

Fuller breasts without surgery and implants

Safe, low-invasive method of breast enlargement without the use of implants, a global breakthrough in aesthetic medicine. Treatment with the use of filler Macrolane facilitate safe and almost painless breast enlargement without surgery and implants.

The new method is part of a global trend of reducing invasiveness in aesthetic medicine endeavors. It provides an excellent solution for women who dream of fuller shape, but fear the scalpel and painful recovery, or are reluctant to silicone implants. Just like any other medical procedure, there may be side effects, such as thickening and nodules - but they disappear through massage. Contraindications to the procedure relate to women at high risk of breast cancer, the occurrence of cysts in the breast glands and changes in clinical breast ultrasound and mammography. Also autoimmune disease (hypothyroidism), liver and kidney, acute and chronic infections, blood clotting disorders, and found hypersensitivity to hyaluronic acid excludes the possibility of implementing the procedure. Macrolane is a product based on the basis of clear gel containing hyaluronic acid stabilized origin. It is worth noting that hyaluronic acid is a natural substance occurring in the skin, subcutaneous tissue and in tissue and joint fluid. The treatment involves injection into each mammary gland preparation in an amount of about 100-120 ml of local anesthesia or intravenous (short term). Breast volume gains and a nicer shape. Effects are visible almost immediately, but it is worth emphasizing that the breasts look natural, even to touch the material in the breast is not perceptible. The procedure is simple, relatively brief and almost painless, and the woman does not need to wear the clothes of any medical specialty. Immediately after the surgery can go home and their daily routine. You should only expect a slight redness and swelling after but these symptoms are evidence of the proper operation of the preparation and subside after 1-2 weeks. This effect persists for about 2 years (depending on the state of the breast, as well as age, style life and physical activity the patient). Until then, hyaluronic acid contained in a preparation is gradually absorbed. The procedure can be repeated after this period, appropriate amount of filler. "The treatment using Macrolane provides the best effect on breast augmentation with cups A to B and the correction of asymmetrical breasts. The method is sensitive, and at the same effect. Does not require post-operative procedure, and can be repeated.