

Cosmetic surgery face and neck

The most common cause of plastic surgery procedures used in these places are very visible changes in birth, accident, or simply post-operative changes resulting from the passage of time.

In this article I will focus on the latter cause. Aesthetic surgery treatments may affect different parts of the face and be due to various reasons. Quite often they are wrinkles, flaccidity of the skin and there are treatments caused a psychological factor. The most common surgery performed on the face and neck treatments facelift (face-lifting), which involves removing excess skin and after then cutting around pre-. Most reviews of plastic surgery offices are women and men after 40 years of age who undergo the treatment in general or local anesthesia (intravenous sedation). The effect of this treatment is less sagging skin and a smaller number of wrinkles. After treatment, the skin in the area cut may be swollen, bruised and painful to the touch, and thus recommends a reduction in physical activity and healing can take several weeks. Condition for carrying out the surgery is a valid general health and normal laboratory results. Another common treatment is a Fine Arts / Eyelid surgery aimed at the same as in the case facelift to remove excess skin, this time around the upper and lower eyelids, wrinkles and excess fat. The procedure is carried out in general or local anesthesia. After the surgery can be problems with Soft Close Automatic for the eyelids, is due to swelling, complete healing occurs within a few weeks. Contraindications to retinal detachment, diabetes, thyroid, glaucoma, dry eye syndrome, hypertension, poor general health, and abnormal results of laboratory tests. Another reason for worry is the shape of many patients and their ears against the head size. The procedure is performed under local anesthesia during which the cut portion of skin and cartilage to be properly arranged. Cutting is performed in an invisible part of the ear which causes the scars are visible. After treatment, lasting an hour your ears should be protected from injury (Sleeping with the band) and avoid low temperatures.